

## THE RESEARCH REGARDING THE PRIMARY COMPACTION OF THE SOILS FROM NORTH WESTERN ROMANIA

Șandor Maria\*, Colibaș Maria

\*University of Oradea, Faculty of Environmental Protection, 26 Gen. Magheru, St. 410048 Oradea; Romania; e-mail: sdaoradea@yahoo.com

### Abstract

*The paper presents a brief summary of research carried out in Sânmartin and Oradea research fields of Agricultural Research and Development Station Oradea.*

*The research from Sânmartin shows the positive effect of the scarification on the physical and chemical properties of soil, increased of the capacity of water storage and of the roots depth and finally yields increases from 25% to 16% in maize and wheat; several series of experiments have revealed a four years duration of the scarification.*

*Research from Oradea were performed in the crop rotation wheat-maize fertilized with  $N_{60}P_{45}$  and in the crop rotation wheat + clover – clover-maize-soybean-maize-oil flax (sunflower) +  $N_{120}P_{90}$  + 50 t/hectares manure shows that scarification caused decreasing of the bulk density values and of the penetration resistance and also increasing of the hydraulic conductivity; the most favourable values of physical parameters of the soil have been registered in the variant with meliorative crop rotation with clover, manure and scarification.*

*Accessible mineral nitrogen content ( $N-NO_3-NH_4$ ), mobile phosphorus and potassium has grown in the two variants with deep loosening. As a result, average on 6 years, in high agrotechnics variants the wheat production has increased with 30%, while that of maize with 12%.*

**Key words:** deep loosening, crop rotation, mineral fertilizers, manure, yield.

### INTRODUCTION

In the North Western Romania, excessive moisture was and can be is the largest spread which is often associated with the advanced argillization and soil compaction (the primary from Bt horizons of illuvial clay soils and secondary anthropogenic which affect arable layer and the one immediately below), the acidity or alkalinity highlighting the need for extensive areas application of the agro-, pedo-, and hydromeliorative works (Domuța C., et al, 2013).

The primary soils compaction from North-Western Romania was studied in the Pedology and Land Reclamation Laboratory of Agricultural Research and Development Station Oradea. In 1968, Colibaș I., introduced for the first time in the Romanian agriculture, tilling with scarifier called "Brăila Progress" worn on the tractor-1300. It was first used in the research field of Sânmartin (Colibas I., 1979).

Use of scarification and subsoiling to control primary soil compaction was also studied by Zahan P., at Husasău Tinca and Oradea. (Şandor M., 2007). This paper presents the research carried out at Sânmartin and Oradea.

## **MATERIAL AND METHOD**

Research from Oradea and Sânmartin carried out in similar climatic conditions: annual rainfall (average for the period 1931-2011) 620,0 mm, the average temperature of the year 10,5<sup>0</sup>C, air humidity 78%. Properties of soil in the two research fields are presented in table 1.

### *Properties of soil from Sânmartin, Bihor research field*

Colloidal clay content reaches at 43% in horizon BtW, textural differentiation is strong (IDT = 1.76), higher bulk density just below the depth of ploughing layer throughout the profile, settlement degree indicates a strong and excessive compaction, total porosity and hydraulic conductivity have small values (Colibas I., 1990).

These properties indicate the absolute necessity of deep lossening of soil. The soil reaction is strongly acid and saturation in the bases degree characterizing the soil as oligo-meso-basic and indicate the necessity for amendmets. Supply with nutrients is medium, so it is necessary to apply fertilizers (Sandor M., 2001).

### *Properties of soil from Oradea research*

Colloidal clay content up to 39%, while physical clay is up to 54%, with a moderate textural differentiation. The settlement degree indicates a weak to strong compaction on profile. Bulk density is middle only in the ploughing layer, high and very high then, and the total porosity and aeration values are medium at surface, then smaller and very smaller. Hydraulic conductivity is good in upper horizons and presents small and very small values below 50 cm deep. Acid soil reaction in Ap horizon passes up in weak acid level till depth, and the contents of the mobile aluminum may disturb development of some crops (Sandor M., 2004).

Table 1

The main properties of soil from research fields

Location	Horizon	Depth cm	Clay <0,002%	BD g/cm <sup>3</sup>	TP %	FC %	K mm/h	pH H <sub>2</sub> O	mobile Al mg/100 g sol	V %	Humus %	P	K
												ppm	
ORADEA	Ap	0-24	31.5	1.33	51	23.0	21.0	5.5	3.7	68	2.32	22.0	83.0
	Et	24-34	31.6	1.38	49	23.2	9.0	5.6	2.3	65	2.28	23.0	102.0
	Bt 1	34-54	40.0	1.44	47	23.7	6.0	6.2	0.5	72	1.91	6.0	112.0
	Bt 2	54-78	39.3	1.55	43	24.6	1.0	6.3	0.8	84	1.93	6.0	118.0
	Bt/C	78-95	39.2	1.62	40	24.6	0.5	6.6	0.3	82	-	-	-
	C	95-145	37.6	1.66	39	24.5	0.1	6.5	0.6	78	-	-	-
SANMARTIN	Ap	0-22	24.7	1.24	53	23.0	5.5	5.1	1.0	51.2	2.80	12.6	59.8
	Eaw	22-32	23.7	1.52	43	23.0	1.9	5.2	1.4	57.6	1.27	11.8	60.0
	EbtW	32-44	30.0	1.54	43	23.0	1.3	5.2	5.5	66.0	1.50	7.8	73.0
	BtW	44-90	43.3	1.63	40	24.0	0.2	5.4	2.4	72.2	-	7.0	63.9
	Btw	90-100	40.2	1.65	39	23.0	0.2	5.9	1.2	81.0	-	-	-

BD – bulk density; TP- total porosity; FC – field capacity; K – hydraulic conductivity.

## RESULTS AND DISCUSSIONS

Research on the effect of deep loosening works through scarification took place in three cycles: in the period 1968 - 1975 it was established the effect and the effect duration of scarification, influence on soil changes and on production increasing; 1975-1979 was researched the effect of scarification by returning on the same ground with scarification works at different time intervals; 1979-1987 were researched aspects of the intensity of the soil scarification, integration of scarification under crop rotation, as well as stabilisation measures for the extension of the effect duration of scarification through pedomeliorative fertilization (on loosening depth) (Colibas I., 1990).

During the period 1974-1987, have initiated the first researches on enhancing the effectiveness of the underground drainage systems through heavy and tamping soils affected by excessive moisture using scarification performed perpendicular to the posing direction of drains to enable the water infiltration into the drains (Colibas I., 1988). Research for primary compaction control and adjusting of aerohidric regime by deep loosening with soil layers inversion without inversion, it began in 1978 on gleyic albic luvisol of Sânmartin- Bihor (Colibas I., 1979).

### **Results obtained in Sânmartin research field**

#### **Effect and duration of soil scarification works**

The biggest physical changes have occurred mainly between 30-60 cm depth: bulk density values have fallen by up to 12%, total porosity increased by up to 15%, porosity aeration values have doubled, and the humidity increased in depth (45-60 cm) with 8%, which explains the large increases obtained at maize crop, which during the summer drought could benefit from a significant water reserve.

Chemical properties have changed under the influence of the 5 t/acres of  $\text{CaCO}_3$  application and of the deep loosening through scarification. Thus, the soil reaction has been obviously improved compared to original state, particularly in the upper soil layers with influences up to 60 cm depth in the case of scarification.

Analysis of mobile nutrient content highlights the fact that the same annual doses of applied fertilizers scarification affect a higher content of nutrients, but it also achieves significant yield increases, as result of the creation of more favorable conditions for the mobilization of soil reserves. Increase proportion of nitric nitrogen compared with the ammoniacal, and the values of the manganese and the ratio of iron as ferric and ferrous highlight the creation of favorable conditions of oxidation processes.

The production capacity of the soil increases being performed significant increases of 25% at corn and of 16% at wheat, the average of the first four years (Sandor M., 2001).

The greatest economical efficiency of scarification was obtained at the maize crop. At wheat crop is also get through scarification, good economical efficiency. In several series of experiments performed at Sânmartin decided that the duration effect of scarification work is 4-5 years, period after which it is necessary to return to the deep loosening soil works.

#### **Effect of soil scarification periodicity**

On the periodicity of the soil scarification have made research on returning with scarification works on the same field yearly, at 2 years, 4 years and 8 years. Both, the results of physical, hydrophysical and micromorphological changes, as well as the level of yields obtained, highlight the fact that the soil scarification execution at intervals of less than 4 years as well as 8 years, is not justified.

#### *Physical and hydrophysical soil changes*

Scarification in different time intervals affects the important changes of the bulk density, penetration resistance and hydraulic conductivity. Thus, bulk density is significantly reduced specially between 20-70 cm depth, depending on the number of deep loosening. The biggest abatements are recorded at every 4 years scarification.

Penetration resistance register declines up to 15%, scarification effect at 2 and 4 years being approached. Hydraulic conductivity increases very significantly under the influence of scarification compared with the soil which is not scarified.

Enhancing the values of this index is being felt throughout the all investigated profile depth, the maximum being at every 4 years scarification (Colibas I., 1990).

#### **Effect of soil scarification in different conditions of agrotehnic level**

Aspects of the scarification effect in different conditions of crop rotation and fertilization were investigated in Sânmartin and Oradea. At Sânmartin we rend the results of a series of experiments with a two-year crop rotation (wheat-maize) and with the four year crop rotation (wheat-clover-corn-flax), at the same level of fertilization ( $N_{120}P_{90}$ ) (table 2).

Yield results, average in the period 1978-1984, obtained from wheat and corn highlight the good effect of soil scarification under the two crop rotations. Yield increases made by scarification are of 360 kg/ha of wheat and respectively of 11% in four year crop rotation and of 400 kg/ha (14%) in two year crop rotation. Crop rotation factor achieves also significant increases of 380 kg/ha wheat (12%) (Colibas I., 1990).

*Table 2*

Influence of scarification and crop rotation on wheat and maize yield in the conditions from Sânmartin-Bihor

Soil  scarification (A)	Crop rotation						Crop rotation average		
	two years			four years (with clover)					
	kg/ha	%	dif	kg/ha	%	dif	kg/ha	%	dif
<b>WINTER WHEAT</b>									
Unscarified	2,900	100	-	3,380	100	-	3,140	100	-
Scarified	3,310	114	400	3,740	111	360	3,520	112	380
<b>MAIZE</b>									
Unscarified	4,260	100	-	4,860	100	-	4,560	100	-
Scarified	4,980	117	720	5,430	112	570	5,200	114	640

For maize, scarification significantly influenced the improve yield with 570 kg/ha (12%) in four year crop rotation and with 720 kg/ha (17% in crop rotation for two years). Rotation factor achieved significant increase of 640 kg/ha (14%). At Oradea in the period 1982-1987 was researched the effect of soil scarification at two levels. Agrotechnical average level included a two year crop rotation (wheat-maize) and annual fertilization N<sub>60</sub>P<sub>45</sub> kg/ha, while the upper level is a six year crop rotation (wheat-clover-maize-soybean-maize-flax oil) with N<sub>120</sub>P<sub>90</sub> fertilization and manure-50 t/ha after maize.

#### Results obtained in theresearch field from Oradea

Physical and hydrophysical soil changes (table 3). Bulk density (BD) decreases by 2-5% depending on the agrotechnical level, particularly on the soil layers from 20 to 40 cm long and 4-8% depending on scarification.

*Table 3*

The influence of agrotechnical level on physical indices from Oradea

Variant	Depth cm	BD g/cm <sup>3</sup>	PR kg/ cm <sup>3</sup>	K mm/h
Average agrotechnics winterwheat – maize*	0-10	1.30	15.8	8.1
	10-20	1.36	15.9	8.7
	20-30	1.53	27.4	5.4
	30-40	1.56	35.2	8.7
Average agrotechnics + scarification winter weath-maize+deep loosening *	0-10	1,28	14.7	9.7
	10-20	1.33	15.0	8.9
	20-30	1.51	25.7	7.6
	30-40	1.49	35.4	8.8
wheat+clover-clover-maize-soybean- maize- flax oil (sun flower) **	0-10	1.26	12.1	8.7
	10-20	1.35	15.6	9.2
	20-30	1.50	27.2	7.6
	30-40	1.49	32.3	8.7
wheat+clover-clover-maize-soybean- maize-flax oil (sun flower) + deep loosening**	0-10	1.23	10.9	12.8
	10-20	1.31	13.3	9.3
	20-30	1.43	28.0	8.0
	30-40	1.44	21.4	8.9

\*- fertilized with N<sub>60</sub>P<sub>45</sub>

\*\* - fertilized with N<sub>120</sub>P<sub>90</sub> + manure, 50 t/ha

Penetration resistance (PR) present values less than 2- 24% in favor of higher agrotechnics compared to average agrotechnics. The scarification influence the reductions of soil penetration resistance by up to 15% compared with the higher agrotechnics. Hydraulic conductivity (K) present low values on soil layer at 20-30 cm, so the presence of hardpan. The scarification enhances this index and in this tamping layer with 41% from the agrotechnics average level and with 48% at the highest. Hydraulic conductivity values grow on all soil layers, from the average level of agrotechnics to the higher and from the unscarified soil to the scarified, with up to 58% (Sandor M., 2001).

*Modifications of some chemical soil indices*

Soil reaction shows a weak trend of acidification after four years from the application dose of N<sub>120</sub> kg/ha/year at the upper agrotechnics level, pH values being lower than 0.2-0.3 units compared to the average agrotechnics level. This trend is weaker in the case of soil scarification on both agrotechnics levels.

Table 4

The influence of agrotechnical level on scarification and on the chemical indices from Oradea (four year effect)

Variant	Depth cm	pH (H <sub>2</sub> O)	N-NO <sub>3</sub>	N-NH <sub>4</sub>	P	K
			ppm		mobil ppm	
Average agrotechnics winter wheat – maize*	0-10	5.32	12.6	6.1	50.2	88.8
	10-20	5.26	14.4	5.7	55.0	87.2
	20-30	5.31	9.5	5.1	51.9	86.3
	30-40	5.50	13.3	6.2	37.5	78.0
Average agrotechnics + scarification winter weath-maize+deep loosening *	0-10	5.01	13.1	4.7	60.6	83.8
	10-20	5.13	14.8	4.6	46.2	88.8
	20-30	5.17	16.4	6.1	49.3	79.7
	30-40	5.15	17.7	5.1	45.5	76.4
wheat+clover-clover- maize-soybean-maize- flax oil (sun flower) **	0-10	5.06	15.9	5.5	64.1	91.3
	10-20	5.03	16.7	5.0	64.6	97.9
	20-30	5.03	17.7	4.6	65.0	86.3
	30-40	5.23	17.6	4.6	47.0	81.3
wheat+clover-clover- maize-soybean-maize- flax oil (sun flower) +deep loosening**	0-10	5.12	15.9	5.3	39.7	88.8
	10-20	5.07	17.3	6.4	39.3	90.5
	20-30	5.17	16.9	6.6	42.3	83.0
	30-40	5.36	18.1	6.4	48.9	93.8

\*- fertilized with N<sub>60</sub>P<sub>45</sub>

\*\* - fertilized with N<sub>120</sub>P<sub>90</sub> + manure, 50 t/ha

Contents of nitric nitrogen in soil is improving under the influence of the upper agrotechnics and scarification on all the investigated soil layers up to 40 cm depth and contrast in soil layers from 20-40 cm who originally presented a lower content and a higher consolidation degree. Soil scarification contributes besides yield improvement at a more uniform distribution on the profile of the mobile nutrients and a more intense mobilization of the existing soil reserves.

*The influence of the agrotechnical level and scarification on yield*

Average yields on six year increase significantly, with 30% on wheat and with 12% for maize from the medium to higher agrotechnical level (table 5). The scarification influence the significant yield achievement of 11- 16% (510-810 kg/ha) to wheat and 12-16% (560-780 kg/ha) for maize, on the two agrotechnical levels (Colibas I., 1990).

*Table 5*  
The influence of scarification and agrotechnical level on yield from soil of Oradea (1982-1987)

Soil scarification (A)	Agrotechnical level III						Average level		
	Crop rotation 2 years + N <sub>60</sub> P <sub>45</sub>			Crop rotation 6 years (with clover N <sub>120</sub> P <sub>90</sub> +50t manure)			Agrotechnical		
	kg/ha	%	dif	kg/ha	%	dif	kg/ha	%	Dif
<b>WHEAT</b>									
Unscarified	3,880	100	-	5,050	100	-	4,460	100	-
Scarified	4,390	111	510	5,860	116	810	5,120	115	660
Scarification average	4,130	100	X	5,460	132	1,330	X	x	X
<b>MAIZE</b>									
Unscarified	4,450	100	-	5,010	100	-	4,730	100	-
Scarified	5,010	112	560	5,790	116	780	5,400	114	640
Scarification average	4,730	100	x	5,400	114	670	X	x	
LSD 5% (A)	630								
LSD 5% (B)	530								

## CONCLUSIONS

North-Western Romania has large areas with secondary compaction. Research concerning the primary compaction control was carried out in the Pedology and Land Reclamation Laboratory of the current Agricultural Research and Development Station Oradea. Research was carried out since 1968 at Sânmartin and Oradea.

In the research field from Sânmartin Colibaş I. was the first researcher from Romania who used the scarification for primary compaction. The results research from Sanmartin emphasized improvement of physical and chemical soil properties mainly to the depth of 30-60 cm in the variants with scarification. The yield increased with 25% in maize and 16% in winter wheat.

The research carried out Oradea in crop rotation wheat-maize fertilized with N<sub>60</sub>P<sub>45</sub> and wheat+clover-clover-maize-soybean-maize-flax oil (sunflower) + N<sub>120</sub>P<sub>90</sub> + 50 t manure/ha shows that scarification determined the decrease of the bulk density and of the penetration resistance values and the increase of the hydraulic conductivity; the most favourable values of soil physical parameters have been registered in the variant with meliorativ crop rotation with clover, manure and scarification.



Accessible mineral nitrogen content ( $N-NO_3-NH_4$ ), mobile phosphorus and potassium have grown in the two variants with deep loosening.

As a result, average on 6 years, in the variant with higher agrotechnics, wheat yield increased with 30% in winter wheat and with 12% in maize.

Research results reflect the importance of deep loosening in the control of the primary soils compaction and the need of the high agrotechnics level (crop rotation, chemical fertilization, organic fertilization).

## REFERENCES

1. Colibaş I., Catargiu D., Canarache A., 1979, Afânarea adâncă a solurilor podzolice şi podzolate cu exces de umiditate temporar. Necesitate şi eficienţă. Rev. Prod, vegetala - cereale şi plante tehnice, nr. 12, pp. 30 – 39.
2. Colibaş I., Mate Şt., 1972, Efectul unor măsuri ameliorative în sporirea producţiei pe solurile podzolice cu exces de umiditate din Bihor. "Zece ani de activitate în sprijinul producţiei", S.E.A. Oradea.
3. Colibaş I., Canarache A., Colibaş Maria, Şandor, Maria, 1990, Efectul lucrărilor de scarificare în asociere cu culturi amelioratoare în asolament, în condiţiile solurilor grele şi tasate din Câmpia piemontană a Crişurilor. Prod. Veg. Cereale şi plante tehnice, nr.2.
4. Colibaş I., Colibaş Maria, 1984, A Roman Szocialista Koztarsasag Eielenyes-Bihar medenceben talalhato idoszakos belvizes talajok melwracios kutatasi eredmenyei. Simpozium "Tessedik Samule", Szarvas, R. P. Ungară.
5. Colibaş I., Colibaş Maria, Şandor Maria, 1988, Cercetări privind cunoaşterea şi ameliorarea unor factori negativi ai fertilităţii solurilor grale şi tasate, afectate de exces de umiditate, din Câmpia Crişurilor şi depresiunile Holod şi Beiuş, în volumul „Contribuţii ale cercetării ştiinţifice la dezvoltarea agriculturii din zona centrală a Câmpiei de vest, 25 de ani de activitate a S.C.A.Z. Oradea”, Bucureşti, Red. de propagandă tehnică agricolă, pp. 445 – 496.
6. Colibaş I., Colibaş Maria, 1983, Rezultate ale cercetărilor privind ameliorarea prin măsuri agro, pedo şi hidroameliorative a solurilor cu exces temporar de umiditate din depresiunea Beiuşului-Bihor. Lucr. Conf. Naţ. pt. Ştiinţa Solului, Brăila, vol. XXI –A, pp. 102 - 109.
7. Colibaş I., Colibaş Maria, 1986, A melylazitas es javitotragyazas hatasa pseudoglejes podzolitajokon. Studia Universitatis Scientiarum Agriculturae, Debreceniensis
8. Colibaş I., Colibaş Maria, Stanga N., 1979, Efectul rescarificării asupra unor indici pedoameliorativi şi recoltei în condiţiile solurilor podzolice argiloiluviale pseudogleice din câmpia piemontana a Crişurilor. Lucr. Conf. Naţ. de Şt. Solului, Braşov, pp. 85 – 91
9. Colibaş I., Colibaş Maria, Şandor Maria, 1995, Cercetări privind sporirea potenţialului productiv al unor lăcovişti din Câmpia joasă a Crişurilor, prin măsuri agro., pedo. şi hidroameliorative. Analele Univ. Oradea, vol I.
10. Colibaş I., Colibaş Maria, Şandor Maria, 1995, Cercetări privind stabilirea unor măsuri agropedoameliorative de valorificare superioară a soloneţurilor din câmpul de drenaj Diosig. Ses. ştiinţ. a SCAZ Oradea.
11. Colibaş I., Colibaş Maria, Şandor Maria, Sabău N.C., 1990, Rezultate ale cercetărilor privind caracterizarea pedologică şi ameliorarea solurilor grele şi tasate, afectate de

- exces de umiditate din zona de activitate a SCAZ Oradea. Conf. Naț. Agrofitehnie, Oradea.
12. Colibaș I., Maria Colibaș, 1988, Rezultate ale cercetărilor privind ameliorarea solurilor cu exces de umiditate din județul Bihor. Publ. S.N.R.S.S. nr. 24, pp. 235 – 259.
  13. Colibaș I., Maria Colibaș, Tatiana Postolache, P. Papacostea, Maria Sandor, 1987, Efectul compactării secundare și a unor măsuri de refacere a stării de afânare, asupra însușirilor solului și producției în condițiile luvisolului pseudogleizat de la Sânmartin - Oradea. Lucr. Conf. Naț. de Șt. Solului, Timișoara, Publ. S.N.R.S.S. nr. 23 A. pp. 129 – 139, București.
  14. Colibaș I., N. Stânga, I. Stanciu, Maria Colibaș, C. Domuța, 1987, Rezultate ale unor cercetări privind ameliorarea solurilor grele și tasate afectate succesiv de exces și deficit de umiditate din Câmpia Crișurilor. Lucr. Conf. Naț. de Știința Solului, Timisoara, Publ. S.N.R.S.S. nr. 23 A, București, pp. 173 - 181.
  15. Colibaș I., și colab., 1985, Veriinderung von hydrophysikalischen Bodeneigenschaften und der Produktion unter dem Einfluss der wiederholten Tieflockerung, Tag-Ber. Akad. Landwirtschaft. Wiss. D.D.R., Berlin, 231.
  16. Colibaș Iuliu, Canarache Andrei, Colibaș Maria, Șandor Maria, 1990, Efectul lucrărilor de scarificare în asocieri cu culturi ameliorative în asolament în condițiile solurilor grele tasate din Câmpia piemontană a Crișurilor. Rev. Producția vegetală – cereale și plante tehnice, nr. 2.
  17. Colibaș, I., Colibaș, Maria, 1983, Efectul drenajului și a unor măsuri pedoameliorative pe luvisolurile albice pseudogleice din depresiunea Beiuș-Bihor, Analele I.C.I.T.I.D., vol. III (XIV).
  18. Șandor Maria, 2004, The influence of the king explotation on the vegetation from canals and of the drainage paramethers from the research field Cefa, Bihor. University of Oradea, Faculty of Enviromment Protection- University of Debrecen. Natural Ressources and Sustainable Development, 23-24 aprilie pag 46.
  19. Șandor Maria, 2007, Ameliorarea solurilor cu exces de umiditate din Câmpia Crișurilor. Editura Universității din Oradea.
  20. Șandor Maria, 2007, Combaterea excesului de umiditate în Câmpia Crișurilor. Editura Universității din Oradea.
  21. Șandor Maria, Dumitru Elisabeta, Ciobanu Gh., Domuța C., Sabău N.C., 2001, Modifications of some physical propieties of the luvic brown soil from Oradea under the technology system influences . Tartamkiserletek. Tajtermesztes, Videkfejlesztés, Nemzetközi Konferencia – Proceedings, Nyiregyhaza-Debrecen, Hungary, pag 221 – 229.
  22. Șandor Maria, și colab., 2004, Physical and chemical properties of the sediments from drains from research field Cefa, Bihor County - Proceeding of University of Debrecen, ISBN 963-472-729-8 HU; ISBN 973-643-379-6 RO, pag.45-46.